

NEW YORK UNIVERSITY

INSTITUTE OF MATHEMATICAL SCIENCES
LIBRARY

AFOSR-TN-60-322

25 Waverly Place, New York 3, N. Y.



NEW YORK UNIVERSITY

Institute of Mathematical Sciences

Division of Electromagnetic Research

RESEARCH REPORT No. BR-32

A Special Hill's Equation with Discontinuous Coefficients

HARRY HOCHSTADT

Mathematics Division

Air Force Office of Scientific Research

Contract No. AF 49(638)-229

Project No. 47500

MARCH, 1960

BR-32
C.1

NEW YORK UNIVERSITY
Institute of Mathematical Sciences
Division of Electromagnetic Research
Research Report No. BR-32
March 1960

A Special Hill's Equation
with Discontinuous Coefficients

Harry Hochstadt

Qualified requestors may obtain copies of this report from the Armed Services Technical Information Agency, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services, or have their 'need to know' certified by the cognizant military agency of their project.

Harry Hochstadt
Harry Hochstadt
Morris Kline
M. Kline
Project Director

Abstract

The equation

$$y'' + \lambda^2 \rho(x)y = 0$$

$$\begin{aligned} \rho(x) &= 1 & |x| &\leq 1 \\ &= a^2 & 1 < |x| < L \end{aligned}$$

is investigated. $\rho(x)$ is taken as a periodic function of period $2L$ and defined as above in the interval $(-L, L)$.

The nature of the characteristic values λ for which the equation has solutions of period L and $2L$ is investigated. Stability charts are drawn showing for which values of a^2 and λ all solutions are bounded, and for which some solutions are not bounded.

Table of Contents

	Page
1. Introduction.	1
2. Preliminary Remarks.	1
3. The Calculation of Characteristic Values for $a^2 > 0$.	4
4. The Stability Chart for $a^2 > 0$.	10
5. The Calculation of Characteristic Values for $a^2 < 0$.	12
6. Applications	16
References	17

1. Introduction.

A particular case of Hill's equation will be treated in detail, namely

$$y'' + \lambda^2 Q(x)y = 0.$$

Here $Q(x)$ will be taken as an even periodic function of period $2L$ and defined by

$$\begin{aligned} Q(x) &= 1 & 0 \leq x < 1 \\ &= a^2 & 1 \leq x \leq L. \end{aligned}$$

Evidently $Q(x)$ is a piecewise constant function with two step discontinuities in the period interval.

2. Preliminary Remarks.

We will first quote some standard results in the theory of Hill's equation. Proofs and references may be found in [1] and [2].

When $Q(x)$ is piecewise continuous, even and has minimal period $2L$, we can find normalized solutions with initial values

$$\begin{aligned} y_1(0) &= 1 & , & & y_1'(0) &= 0 & , \\ y_2(0) &= 0 & , & & y_2'(0) &= 1 & . \end{aligned}$$

The solutions of the characteristic equation

$$\rho^2 - 2y_1(2L)\rho + 1 = 0$$

are known as characteristic exponents. According to Floquet's theorem, we

can then find solutions $f_1(x)$, $f_2(x)$ such that

$$f_1(x + 2L) = \rho_1 f_1(x)$$

$$f_2(x + 2L) = \rho_2 f_2(x)$$

when ρ_1 and ρ_2 are distinct characteristic exponents. If $\rho_1 = \rho_2$, i.e.

$\rho_1 = \rho_2 = 1$ or $\rho_1 = \rho_2 = -1$, we can find a solution $y(x)$ with the property

$$y(x + 2L) = \rho y(x).$$

If $\rho = 1$, $y(x)$ has period $2L$; if $\rho = -1$, $y(x)$ has period $4L$.

Furthermore, we have a stability test. Solutions are stable if and only if

$$|y_1(2L)| < 1$$

$$\text{or} \quad y_1(2L) = \pm 1 \text{ and } y_2(2L) = y_1'(2L) = 0.$$

Evidently, for $|y_1(2L)| > 1$, ρ_1 and ρ_2 are real and since $\rho_1 \rho_2 = 1$ at least one of them is greater than one in absolute value. The solution corresponding to that exponent evidently has increasing peak values.

Another important statement is the following: There exists a non-trivial periodic solution which is

- 1.) even and of period $2L$ if and only if $y_1'(L) = 0$,
- 2.) odd and of period $2L$ if and only if $y_2(L) = 0$,
- 3.) even and of period $4L$ if and only if $y_1(L) = 0$,
- 4.) odd and of period $4L$ if and only if $y_2'(L) = 0$.

By means of the Liouville transformation [3] we can transform equations of the form

$$f_2(x)y''(x) + f_1(x)y'(x) + [\lambda g(x) + f_0(x)]y(x) = 0$$

into the form

$$\eta''(\xi) + [\lambda + \psi(\xi)]\eta(\xi) = 0.$$

In order to effect this transformation certain differentiability conditions must be imposed on the coefficients in the equation. In particular, to apply the Liouville transformation to an equation of the type

$$y'' + \lambda^2 Q(x)y = 0$$

it is necessary that $Q(x)$ possess a second derivative. In the particular case under discussion these conditions are not fulfilled. A certain Oscillation Theorem has been proved by Haupt [2], which states that the equation

$$\eta''(\xi) + [\lambda + \psi(\xi)]\eta(\xi) = 0,$$

where $\psi(\xi)$ is a doubly differentiable, real function of period $2L$, has solutions of period $2L$ for an infinite sequence of characteristic values $\lambda_0 < \lambda_1 < \lambda_2 < \dots$, and solutions of period $4L$ for a sequence $\lambda'_1 < \lambda'_2 < \lambda'_3 < \dots$. These λ 's satisfy the inequalities

$$\lambda_0 < \lambda'_1 \leq \lambda'_2 < \lambda_1 \leq \lambda_2 < \lambda'_3 \leq \lambda'_4 < \lambda_3 \leq \lambda_4 < \dots$$

and satisfy the relations

$$\lim_{n \rightarrow \infty} \lambda_n^{-1} = 0, \text{ and}$$

$$\lim_{n \rightarrow \infty} \lambda'_n^{-1} = 0.$$

Furthermore, for λ 's in the intervals (λ_0, λ'_1) , (λ'_2, λ_1) , (λ_2, λ'_3) , (λ'_4, λ_3) , ... the solutions are stable, outside of these intervals unstable and at the end points they are in general unstable, except under very special conditions.

We will investigate to what extent these theorems remain valid for the equation under discussion.

3. The Calculation of the Characteristic Values for $a^2 > 0$.

We now turn to the equation

$$y''(x) + \lambda^2 y(x) = 0, \quad 0 \leq x < 1$$

$$y''(x) + \lambda^2 a^2 y(x) = 0, \quad 1 \leq x \leq L$$

and investigate the characteristic values under the boundary conditions

$$1.) \quad y'(0) = y'(L) = 0 \quad (\text{even solutions of period } 2L)$$

$$2.) \quad y(0) = y(L) = 0 \quad (\text{odd solutions of period } 2L)$$

$$3.) \quad y'(0) = y(L) = 0 \quad (\text{even solutions of period } 4L)$$

$$4.) \quad y(0) = y'(L) = 0 \quad (\text{odd solutions of period } 4L).$$

Corresponding to the first case, one can show by a direct calculation that

$$\begin{aligned} y(x) &= \cos \lambda x, \quad 0 \leq x \leq 1 \\ &= \cos \lambda \cos \lambda a(x - 1) - \frac{1}{a} \sin \lambda \sin \lambda a(x - 1), \quad 1 \leq x \leq L. \end{aligned}$$

These solutions obviously satisfy the condition $y'(0) = 0$, and from $y'(L) = 0$ we obtain, immediately, the equation

$$a \cos \lambda \sin \lambda a(L - 1) + \sin \lambda \cos \lambda a(L - 1) = 0.$$

An equivalent form is

$$\tan \lambda = -a \tan \lambda a(L - 1).$$

In a similar manner we can construct corresponding equations for the other three cases and can write down the following set of characteristic equations:

$$1.) \quad \tan \lambda = -a \tan \lambda a(L-1)$$

$$2.) \quad \tan \lambda = -\frac{1}{a} \tan \lambda a(L-1)$$

$$3.) \quad \tan \lambda = a \cot \lambda a(L-1)$$

$$4.) \quad \tan \lambda = \frac{1}{a} \cot \lambda a(L-1).$$

We could arrive at the characteristic values of λ by starting with the characteristic exponents defined by

$$\rho^2 - 2y_1(2L)\rho + 1 = 0,$$

and asking for the values of λ for which

$$y_1(2L) = 1 \quad \text{and} \quad y_1(2L) = -1.$$

The first of these refers to the cases labeled 1.) and 2.) above and the second to the cases 3.) and 4.). A direct calculation shows that

$$y_1(2L) = \cos 2\lambda a(L-1) \cos 2\lambda - \frac{1}{2}\left(a + \frac{1}{a}\right) \sin 2\lambda \sin 2\lambda a(L-1)$$

and we can see that

$$\begin{aligned} y_1(2L) - 1 = & -2 \cos^2 \lambda \cos^2 \lambda a(L-1) [\tan \lambda + a \tan \lambda a(L-1)] \\ & \times \left[\tan \lambda + \frac{1}{a} \tan \lambda a(L-1) \right] \end{aligned}$$

showing that the roots of

$$y_1(2L) - 1 = 0$$

correspond to the roots of the characteristic equations of cases 1.) and 2.).

Similarly,

$$y_1(2L) + 1 = 2 \cos^2 \lambda \sin^2 \lambda a(L-1) [\tan \lambda - a \cot \lambda a(L-1)] \\ \times \left[\tan \lambda - \frac{1}{a} \cot \lambda a(L-1) \right]$$

showing the factorization corresponding to cases 3.) and 4.).

The following method will be used to obtain the characteristic values. We treat case 1.) as typical and write the equation in the form

$$\lambda a(L-1) = -\tan^{-1} \frac{1}{a} \tan \lambda.$$

In the evaluation of the inverse tangent we must take the multivalued character of the function into account. We will graph both the right side and the left side of the equation and wherever the curves intersect, we evidently have a characteristic value.

Figure 1 was drawn for $a > 1$. For $a = 1$ the wavy lines become straight and have slope -1. If we superimpose on this figure the other three cases we obtain figure 2.

Figure 2 was drawn for $a > 1$. For $a = 1$ the curves of the type 1.) and 2.) become identical straight lines of slope -1, and similarly for the curves of type 3.) and 4.). Wherever the line $\lambda a(L-1) = f(\lambda)$ intersects these curves, we obtain characteristic values. If $\{\lambda_1\}$ denote the characteristic values corresponding to solutions of period $2L$ and $\{\lambda_1'\}$ the ones corresponding to solutions of period $4L$, we find that

$$0 = \lambda_0 < \lambda_1' \leq \lambda_2' < \lambda_1 \leq \lambda_2 < \lambda_3' \leq \lambda_4' < \lambda_3 \leq \lambda_4 < \dots$$

Thus Haupt's Oscillation Theorem remains valid.




Figure 1




Figure 2

For $a < 1$ the curves of type 1.) and 2.) change place and similarly for types 3.) and 4.).

We can immediately determine where multiple roots occur. Evidently for $a = 1$ we have

$$\left. \begin{aligned} \lambda'_{2i+1} &= \lambda'_{2i+2} \\ \lambda_{2i+1} &= \lambda_{2i+2} \end{aligned} \right\} \quad i = 0, 1, 2, \dots$$

In general even when $a \neq 1$, we have multiple roots for $a(L-1)$ rational, say p/q , where p and q are relatively prime. Then we have double roots at the lattice points (kp, kq) such that for

$$\left. \begin{aligned} k(p+q) &= 2n, & \lambda_{2n-1} &= \lambda_{2n} \\ k(p+q) &= 2n+1, & \lambda'_{2n+1} &= \lambda'_{2n} \end{aligned} \right\} \quad n = 1, 2, \dots$$

It also follows that for $a(L-1) = p/q$, we need to compute only the first p^2 roots of each kind and all others are equal to these modulo $q\pi$.

For $a(L-1) = 1$ we can also see that since

$$\tan \lambda'_{2i+1} = a \cot \lambda'_{2i+1}$$

$$\tan \lambda'_{2i+2} = \frac{1}{a} \cot \lambda'_{2i+2}$$

it follows that

$$\tan^2 \lambda'_{2i+1} \tan^2 \lambda'_{2i+2} = 1$$

so that

$$\lambda'_{2i+1} + \lambda'_{2i+2} \equiv \frac{\pi}{2} \pmod{\pi}.$$

To obtain estimates for the roots we see that for case 1.)

$$\lambda a(L - 1) = -\tan^{-1} \frac{1}{a} \tan \lambda \approx n\pi - \lambda.$$

From this we see that

$$\lambda_{2n-1} = \left[\frac{2n}{1 + a(L - 1)} \right] \frac{\pi}{2} + \theta \frac{\pi}{2}, \quad 0 \leq \theta < 1.$$

(Here $[x]$ represents the greatest integer less than or equal to x .) Similarly,

$$\lambda_{2n} = \left[\frac{2n}{1 + a(L - 1)} \right] \frac{\pi}{2} + \theta \frac{\pi}{2},$$

$$\lambda'_{2n-1} = \left[\frac{2n - 1}{1 + a(L - 1)} \right] \frac{\pi}{2} + \theta \frac{\pi}{2},$$

$$\lambda'_{2n} = \left[\frac{2n - 1}{1 + a(L - 1)} \right] \frac{\pi}{2} + \theta \frac{\pi}{2}.$$

To obtain better numerical estimates in particular cases we can use Newton's method.

4. The Stability Chart for $a^2 > 0$.

We can make a rough stability chart from Fig. 1. That is, we will make a graph showing the relationship between the characteristic λ 's and a . These curves subdivide the first quadrant into regions of stability and instability. That is, for any λ and a from a stable region, all solutions of the differential are stable. Outside the stability regions there is only one bounded solution.




Figure 3

The shaded regions represent regions of stability. The values of λ where the curves of type 2.) and 3.) intersect the abscissa are roots of the equations

$$\tan \lambda = -\lambda(L-1)$$

$$\tan \lambda = \frac{1}{\lambda(L-1)}$$

respectively.

5. The Calculation of Characteristic Values for $a^2 < 0$.

To treat the case where $a^2 < 0$, we return to the previous equations and replace a by ia . Then we have

$$y'' + \lambda^2 y = 0, \quad 0 \leq x < 1$$

$$y'' - \lambda^2 a^2 y = 0, \quad 1 < x \leq L,$$

subject to the boundary conditions

$$1.) \quad y'(0) = y'(L) = 0$$

$$2.) \quad y(0) = y(L) = 0$$

$$3.) \quad y'(0) = y(L) = 0$$

$$4.) \quad y(0) = y'(L) = 0.$$

The corresponding characteristic equations are

$$1.) \quad \tan \lambda = a \tanh \lambda a(L-1)$$

$$2.) \quad \tan \lambda = -\frac{1}{a} \tanh \lambda a(L-1)$$

$$3.) \quad \tan \lambda = a \coth \lambda a(L-1)$$

$$4.) \quad \tan \lambda = -\frac{1}{a} \coth \lambda a(L-1).$$

To evaluate the roots of these equations, we write, as before, in case 1.)

$$\lambda a(L - 1) = \tanh^{-1} \frac{1}{a} \tan \lambda$$

and graph the right side of the equation. We then obtain Figure 4, where all four cases are included.




Figure 4

For $a^2(L - 1) < 1$ we again have a sequence of characteristic roots in the order given by the Oscillation Theorem

$$0 = \lambda_0 < \lambda'_1 < \lambda'_2 < \lambda_1 < \lambda_2 < \lambda'_3 < \lambda'_4 < \lambda_3 < \lambda_4 < \dots$$

In this case we obtain no double roots. For $a^2(L - 1) > 1$ a direct analysis shows that we have two roots corresponding to case 1.) in the interval $[0, \tan^{-1}a]$. Then the sequence becomes

$$0 = \lambda_0^{(1)} < \lambda_0^{(2)} < \lambda'_1 < \lambda'_2 < \lambda_1 < \lambda_2 < \lambda'_3 < \lambda'_4 < \lambda_3 < \lambda_4 < \dots$$

For $a^2(L - 1) = 1$ $\lambda_0^{(1)} = \lambda_0^{(2)} = 0$.

For large n we see readily that

$$\lambda_{2n} \approx n\pi + \tan^{-1}a$$

$$\lambda'_{2n+1} \approx n\pi + \tan^{-1}a$$

$$\lambda'_{2n+2} \approx (n+1)\pi - \tan^{-1} \frac{1}{a}$$

$$\lambda_{2n+1} \approx (n+1)\pi - \tan^{-1} \frac{1}{a}.$$

Figure 5 is the stability chart for this problem, where the regions of stability are shaded.




Figure 5

6. Applications

The equation discussed in Sections 2 and 3 has numerous physical applications. It can be derived from the problem of a vibrating string, composed of two homogeneous pieces. Another application is to the propagation of a wave in a one-dimensional structure, where the index of refraction is an even periodic, piecewise constant function.

Another application is to resonant inductance capacitance circuits, where the inductance or capacitance varies periodically with time. Such equations were treated approximately by Carson ^[4] and Minorsky ^[5].

References

- [1] Coddington, E.A. and Levinson, W. - Theory of Ordinary Differential Equations; New York, McGraw-Hill, 1955.
- [2] Magnus, W. - Hill's equation. Part I: General theory; N.Y.U., Inst. Math. Sci., Div. EM Res., Research Report No. BR-22, June, 1957.
- [3] Kamke, E. - Differentialgleichungen Lösungsmethoden und Lösungen, Band 1. Gewöhnliche Differentialgleichungen. New York.
- [4] Carson, J.R. - Forced and free oscillations in circuits including variable inductance; Internal Memorandum, A.T.T., September 15, 1919. Appendix by C.M. Hebbert, July, 1921.
- [5] Minorsky, N. - On parametric excitation; Franklin Inst. J., 240, 25 (1945).

DISTRIBUTION LIST FOR RESEARCH REPORTS

Contract No. AF 49(638)229

(ONE copy unless otherwise noted)

- (3) Commander
European Office, ARDC
47 Rue Cantersteen
Brussels, Belgium
- Applied Mathematics and Statistics Lab.
Stanford University
Stanford, California
- Department of Mathematics
University of California
Berkeley, California
- Commander
Air Force Flight Test Center
ATTN: Technical Library
Edwards Air Force Base, California
- The Rand Corporation
Technical Library
1700 Main Street
Santa Monica, California
- Director of Advanced Studies
Air Force Office of Scientific Research
Post Office Box 2035
Pasadena 2, California
- Commander
Western Development Division
ATTN: WDSIT
P.O. Box 262
Inglewood, California
- Department of Mathematics
Yale University
New Haven, Connecticut
- Chief of Naval Operations
Department of the Navy
ATTN: OP03EG; Mr. Arthur Kaufman
Washington 25, D.C.
- (2) Office of Naval Research
Department of the Navy
ATTN: Code 432
Washington 25, D.C.
- Department of Commerce
Office of Technical Services
Washington 25, D.C.
- Director National Security Agency
ATTN: Dr. H.H. Campaigne
Washington 25, D.C.
- Library
National Bureau of Standards
Washington 25, D.C.
- National Applied Mathematics Labs.
National Bureau of Standards
Washington 25, D.C.
- Headquarters, USAF
Assistant for Operations Analysis
Deputy Chief of Staff, Operations, AF00A
Washington 25, D.C.
- (2) Commander
Air Force Office of Scientific Research
ATTN: SRDB
Washington 25, D.C.
- (2) Commander
Air Force Office of Scientific Research
ATTN: SRE
Washington 25, D.C.
- National Science Foundation
ATTN: Dr. Leon Cohen
Washington 25, D.C.
- Commander
Air Force Armament Center
ATTN: Technical Library
Eglin Air Force Base, Florida
- Commander
Air Force Missile Test Center
ATTN: Technical Library
Patrick Air Force Base, Florida
- Department of Mathematics
Northwestern University
Evanston, Illinois
- Institute for Air Weapons Research
Museum of Science and Industry
University of Chicago
Chicago 37, Illinois
- Department of Mathematics
University of Chicago
Chicago 37, Illinois
- Department of Mathematics
University of Illinois
Urbana, Illinois
- Department of Mathematics
Purdue University
Lafayette, Indiana
- Institute for Fluid Dynamics
and Applied Mathematics
University of Maryland
College Park, Maryland
- Mathematics and Physics Library
The Johns Hopkins University
Baltimore, Maryland
- Commander
RQ Air Research and Development Command
Director of Research
Andrews Air Force Base
Washington 25, D.C.
- Department of Mathematics
Harvard University
Cambridge 38, Massachusetts
- Department of Mathematics
Massachusetts Institute of Technology
Cambridge, Massachusetts
- Commander
Air Force Cambridge Research Center
ATTN: Geophysics Research Library
L.G. Hanscom Field
Bedford, Massachusetts
- Commander
Air Force Cambridge Research Center
ATTN: Electronics Research Library
L.G. Hanscom Field
Bedford, Massachusetts
- Psychological Labs.
ATTN: Dr. George A. Miller
Memorial Hall
Harvard University
Cambridge 38, Mass.
- Department of Mathematics
Wayne University
ATTN: Dr. Y.W. Chen
Detroit 1, Michigan
- Willow Run Research Center
University of Michigan
Ypsilanti, Michigan
- Department of Mathematics
Folwell Hall
University of Minnesota
Minneapolis, Minnesota
- Department of Mathematics
Institute of Technology
Engineering Building
University of Minnesota
Minneapolis, Minnesota
- Department of Mathematics
Washington University
St. Louis 5, Missouri
- Department of Mathematics
University of Missouri
Columbia, Missouri
- Linda Hall Library
ATTN: Mr. Thomas Gillis
Document Division
5109 Cherry Street
Kansas City 10, Missouri
- Commander
Strategic Air Command
ATTN: Operations Analysis
Offutt Air Force Base
Omaha, Nebraska
- The James Forrestal Res. Center Lib.
Princeton University
Princeton, New Jersey
- Library
Institute for Advanced Study
Princeton, New Jersey
- Department of Mathematics
Fine Hall
Princeton University
Princeton, New Jersey
- Commander
AF Missile Development Center
ATTN: Technical Library
Holloman Air Force Base, New Mexico
- Commander
Air Force Special Weapons Center
ATTN: Technical Library
Kirtland Air Force Base
Albuquerque, New Mexico
- Prof. J. Wolfowitz
Mathematics Department
White Hall
Cornell University
Ithaca, New York
- Department of Mathematics
Syracuse University
Syracuse, New York
- New York University
Institute of Mathematical Sciences
Division of Electromagnetic Research
ATTN: Professor M. Kline
25 Waverly Place
New York 3, New York
- Department of Mathematics
Columbia University
ATTN: Professor B.O. Koopman
New York 27, N.Y.
- Department of Mathematical Statistics
Fayerweather Hall
ATTN: Dr. Herbert Robbins
Columbia University
New York 27, N.Y.
- Commander
Rome Air Development Center
ATTN: Technical Library
Griffies Air Force Base
Rome, New York
- Institute for Aeronautical Sciences
ATTN: Librarian
2 East 61st Street
New York 21, N.Y.

MAR 19 1987		
MAR 20 1987		
SEP 22 1987		
OCT 05 1987		
OCT 17 1987		
NOV 13 1987		
(6)	PRINTED IN U.S.A.	

PRINTED IN U. S. A.

NYU
BR-
32 Hochstadt

c.1

A special Hill's equation
with discontinuous co-
efficients

NYU
BR-32 Hochstadt

c.1

AUTHOR

A special Hill's equation
TITLE with discontinuous
coefficients

DATE DUE

BORROWER'S NAME

ROOM
NUMBER

MAR 13 1961

S. Gorenstein

MAR 29 1979

S. Gorenstein

SEP 22 1997

S. Y. Shem
Shem

N. Y. U. Institute of
Mathematical Sciences
25 Waverly Place
New York 3, N. Y.

